

FOR DIRECT INJECTION INTO DUMPS, HYDRO COOLERS, SPRAY SYSTEM AND PROCESS WATER

General Information

DIRECTIONS FOR USE

SaniDate 5.0 Sanitizer/Disinfectant works best when diluted with water containing minimal levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. This product will readily mix with clean, neutral water and does not require agitation.

SaniDate 5.0 Sanitizer/Disinfectant concentrate should not be combined or mixed with any other pesticide concentrates.

This product is not for use on medical device surfaces.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. A potable water rinse is not allowed (Do not rinse) following use of the product as a sanitizer on previously cleaned hard surfaces provided that the surfaces are adequately drained before contact with food so that little or no residue remains.

Limitations, Restrictions, and Exceptions

For direct injection into dump tanks, hydro coolers, spray system and process waters For treatment of water containing plant pathogens, inject SaniDate 5.0 Sanitizer/Disinfectant and maintain a predetermined residual level by using metering equipment, coupled with ORP measuring probes.

1. Determine biological organic loading prior to treatment if possible.
2. For waters that contain low levels of biological and organic loading inject SaniDate 5.0 Sanitizer/Disinfectant at 1.07 - 0.535 fl. oz. of concentrate for every 100 gallons of water or at a dilution rate of 1:5000 - 10,000.
3. For clean water, inject 0.535 - 0.2675 fl. oz. of SaniDate 5.0 Sanitizer/Disinfectant for every 100 gallons of water or a dilution rate of 1:10,000- 1:20,000 to prevent the

formation of algae, bacteria and fungi.

Method

[Injection](#)

[Spray](#)

Rates

[field rates 0](#)

-

Timings

[Post-harvest](#)