

SURFACE FEEDING LAWN INSECTS

General Information

DIRECTIONS FOR USE

In New York, this product may NOT be applied to any grass or turf area (lawns) within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of Bifenthrin 0.1% Lawn Granules if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Apply when insect activity or damage is first observed in the lawn (lawn thins and turns brown or moths fly from the grass when you mow). Don't treat just the browned area of lawn. To prevent further damage, treat entire lawn.

Limitations, Restrictions, and Exceptions

Uniformly apply 1.2 to 2.4 pounds over a 1,000 sq. ft. area of lawn using any hand or power operated gravity or rotary type spreader designed for the application of granular materials. A [X] pound bag will treat a [X] to [X] sq. ft. area of lawn. Water thoroughly immediately after application. Treat when insects first appear and retreat as necessary. Repeat application is necessary only if there are signs of renewed insect activity. Repeat application should be limited to no more than once per seven days.

Food utensils such as tablespoons/teaspoons must not be used for food purposes after use in measuring pesticides.

Method

[Surface](#)

Rates

[field_rates 0](#)

-

Timings

[When insects first appear and retreat as necessary.](#)