

## **CEREAL GRAINS - BARLEY, BUCKWHEAT, ETC. - ARMYWORM, BIRD CHERRY-OAT APHID, ETC.**

### General Information

#### GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals. per acre by air or 10 gals. per acre by ground unless otherwise specified in the label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, CAVALRY may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

#### RESISTANCE MANAGEMENT

CAVALRY is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

#### Limitations, Restrictions, and Exceptions

#### Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.
- Apply with ground or air equipment using sufficient water and application methods to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre.
- Do not allow livestock to graze in treated areas or harvest treated wheat forage as feed for meat or dairy animals within 7 days after treatment. Do not feed treated straw to meat or dairy animals within 30 days after the last treatment.

Bird Cherry-Oat Aphid, English Grain Aphid, Russian Wheat Aphid, English Grain Aphid, Russian Wheat Aphid - Best control is obtained before insects begin to roll leaves. Once crop has started to boot, CAVALRY may provide suppression only. Higher rates and increased coverage will be necessary.

Hessian Fly, Hessian Fly - Make applications when adults emerge.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

30 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[N.A.](#)