

DRIED SHELLLED LEGUME VEGETABLES (BEANS AND PEAS) - ALFALFA CATERPILLAR, APHID, ARMYWORM, ETC.

General Information

GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals. per acre by air or 10 gals. per acre by ground unless otherwise specified in the label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, CAVALRY may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

RESISTANCE MANAGEMENT

CAVALRY is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Limitations, Restrictions, and Exceptions

Remarks

- Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds.

- Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 2 gals. of water per acre.

- For succulent and dried shelled peas and beans, do not graze livestock in treated areas or harvest vines for forage or hay.

Aphid species, Plant Bug species Including Lygus species, Tobacco Budworm - See Resistance statement under General Directions for Use.

Armyworm, Fall Armyworm, Western Yellowstriped Armyworm, Yellowstriped Armyworm - Use higher rates for large larvae.

Curculio and Weevil species (foliage and pod feeding adults and larvae), Stalk Borer - For control before the larva bores into the plant stalk or pods.

Thrips species - See Resistance statement under General Directions for Use. Does not include Western Flower Thrips.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

24 hours

Timings

[N.A.](#)