

## **NON-CROPLAND (EXCLUDING PUBLIC LAND)**

### General Information

#### GENERAL DIRECTIONS FOR USE

Initial and residual control are contingent upon thorough crop coverage. Apply with ground or air equipment using sufficient water to obtain full coverage of foliage. Apply in a minimum of 2 gals. per acre by air or 10 gals. per acre by ground unless otherwise specified in the label. When foliage is dense or pest pressure is high (heavier insect or egg pressure, larger larval stages), use of higher application volumes and/or higher use rates may improve initial and residual control.

For cutworm control, CAVALRY may be applied before, during, or after planting. For soil incorporated applications, use higher rates for improved control.

#### RESISTANCE MANAGEMENT

CAVALRY is a Group 3 Insecticide (contains the active ingredient lambda-cyhalothrin). Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

#### Limitations, Restrictions, and Exceptions

#### Remarks

- Spray non-cropland adjacent to agricultural areas to control migratory insects,

which may threaten crops.

- Follow General Use Directions, rates and spray recommendations found elsewhere in the label for the adjacent crop outlet and target pests.
- Use highest labeled rates for dense/large foliage, high insect populations and larger larval stages.
- Repeat as necessary to maintain control.
- Do not graze livestock in treated areas.
- Refer the conversion chart below.

Method

[Broadcast/Foliar Air](#)

Rates

[field\\_rates 0](#)

•

Restricted Entry Interval

24 hours

Timings

[N.A.](#)