

CROP RECOMMENDATIONS: VEGETABLE CROPS

General Information

Cytokin is a nutritional supplement derived from sea plant extracts (*Ascophyllum nodosum*) and plant nutrients to supplement the nutritional needs of crops.

APPLICATION INSTRUCTIONS:

For specific crop recommendations, contact your Farm Advisor, PCA, or any Company Representative. Apply with sufficient water to get thorough foliage coverage, 3 to 10 gallons water per acre for aircraft sprayers and 10 to 250 gallons water per acre for ground driven spray equipment. Cytokin may be used with a surfactant.

FOLIAR SPRAY APPLICATION RATE: Apply Cytokin at the rate of $\frac{1}{2}$ to 2 pints per sprayed acre weekly beginning one week after transplanting or at the 3 to 6 leaf stage and continue weekly application up to three weeks before final harvest.

Use this product in accordance with good agronomic practices, which include utilizing proven spray equipment set for proper coverage. Do not make applications when temperatures are too hot. Applications should be made at temperature levels and when other environmental conditions in your area are such that experience indicates the application will accomplish the desired result.

Limitations, Restrictions, and Exceptions

CROP RECOMMENDATIONS: VEGETABLE CROPS

Apply $\frac{1}{2}$ to 2 pints per sprayed acre at one to two week intervals from the 3 to 6 leaf stage until two weeks before harvest.

Method

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

From the 3 to 6 leaf stage.