

FOLIAR SPRAY USE

General Information

GENERAL INFORMATION

FERTIGARD is a nutritional blend designed to be used as a supplement to a balanced fertility program. Foliar applications of FERTIGARD may be beneficial during physiological periods of peak nutrient demand, periods of environmental stress or to crops suffering from a weakened root system. FERTIGARD may also be applied as a post harvest application to improve the vigor of overwintering fruit buds and potential return yields where crop loads were heavy the prior year. Tissue and soil nutrient analysis may be helpful to monitor and maintain plant growth and development.

FOLIAR APPLICATION is intended as a supplement to a regular fertilization program and will not, by itself provide all the nutrients normally required by agricultural crops.

Compatibility: This product is compatible with most pesticides commonly used on agronomic crops. CAUTION: Do not use with hydrated lime, lime sulfur or other highly alkaline materials. FERTIGARD may be applied with copper crop protection materials provided spray solution pH is maintained at 6.5 or higher. Do not apply with silicone based adjuvants or crop oil concentrates.

APPLICATION

FERTIGARD may be applied as a foliar spray by ground, air, concentrate or dilute. For permanent crops, ground dilute treatments are the preferred method of foliar nutrient application.

Refer in the label for mixing directions.

Limitations, Restrictions, and Exceptions

DIRECTIONS FOR USE

Rate of Application in Season (Foliar Spray Use): Use at the rate of 2 to 3 lbs. per acre on all crops, as needed to improve plant vigor and to promote crop production.

Multiple low rate applications are generally more effective than a large single dose. Begin applications early in the growing season and during plant production periods at 10 to 14 days intervals. Post harvest application may be made at 2.5 to 5 lbs. per acre. Applications should be made in a minimum of 20 gallons of water by ground and 5 gallons of water by air.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

•

Timings

[Early in the growing season and during plant production periods.](#)