

## **FOLIAR SPRAYS (FRUIT, VEGETABLE AND ROW CROPS)**

### General Information

Compatibility: This product is compatible with most pesticides commonly used on vegetable, fruit and agronomic crops. CAUTION: Do not use with Dormant Oil, Lime Sulfur, Spray Lime, or highly alkaline spray materials. Add product to the spray tank or pre-mix tank when it is about half full. Agitation in the tank is preferred to increase the speed of solubility. Where extremely cold water is used, allow a few extra minutes for the product to completely dissolve before spraying.

Use this product in accordance with good agronomic practices, which include utilizing proven spray equipment set for proper coverage. Do not make applications when temperatures are too hot, as crop damage may occur. Applications should be made at temperature levels and when other environmental conditions in your area are such that your experience indicates the application will be compatible and will accomplish the desired result.

### Limitations, Restrictions, and Exceptions

#### FOLIAR SPRAYS (FRUIT, VEGETABLE AND ROW CROPS)

To improve early vigor and fruiting, apply SOL-U-GRO MAX in a 10" to 14 foliar banded spray at 2 lbs per acre every 7 to 10 days. Full season foliar sprays may be used at 3 to 10 lbs per acre.

#### Method

[Foliar spray](#)

[Band](#)

Rates

[field rates 0](#)

•

Timings

[N.A.](#)