

CITRUS, POME, STONE AND TROPICAL FRUITS - CONTROL OF MITES

General Information

GENERAL INFORMATION

- Thorough coverage is necessary to provide good disease and mite control. Trilogy prevents fungal attack of plant tissue and contact activity to mite pressure on the plant. Avoid excessive runoff of spray material for maximum control.
- For best results, maintain constant agitation in spray equipment.
- For optimal performance, do not mix with cold water (less than 45°F).
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted crops prior to root establishment.
- Do not apply to known sensitive plant species, such as impatiens flowers, fuchsia flowers, hibiscus flowers, some rose flowers, ornamental olive trees, or some carnation varieties without prior testing.
- When used in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale.
- As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use with care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.

AGRICULTURAL USE DIRECTIONS

Trilogy is a broad spectrum fungicide of certain diseases and controls mites in citrus, deciduous fruits and nuts, vegetable crops, cereal grains, and other miscellaneous crops.

- Do not use Trilogy after bloom on table grapes or following bunch closure on wine grapes. Do not use Trilogy after pit hardening on stone fruit.
- Use care when applying near streams, ponds, lakes, or other bodies of water. Do not apply Trilogy when weather conditions favor drift or when the likelihood of runoff is high.
- Apply Trilogy by conventional, low volume, and aerial application equipment.

Limitations, Restrictions, and Exceptions

CITRUS

- Use Trilogy on citrus to control mites at a rate of 1.0-2.0% in 25-200 gallons of water for high volume application or 0.5 gallon Trilogy in a minimum of 5 gallons of water per acre for low volume application. As a dormant oil spray, use Trilogy as a 2% solution or 1 quart Trilogy minimum per acre.

STONE AND TROPICAL FRUITS

- Use Trilogy at first sign of pests or pest damage at the appropriate rate. Repeat applications as needed for adequate control every 7-21 days. Apply a 1.0-2.0% solution of Trilogy for high volume application or a minimum of 1 quart Trilogy in a minimum of 5 gallons of water per acre for low volume application. Do not use Trilogy after pit hardening on stone fruit.

POME (DECIDUOUS FRUITS) AND NUTS

- Use Trilogy at a rate of 1.0-2.0% in 25-100 gallons of water per acre or 0.5 quarts Trilogy in a minimum of 5 gallons of water per acre for low volume application. Apply no more than 2 gallons of Trilogy per acre per solution. Apply at first sign of pests or pest damage. Repeat application as needed for adequate control every 7-21 days.

Use the dilution tables to calculate dilution rates (please refer to the label).

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

4 hours

Timings

[First sign of pests or pest damage.](#)