

# **CUCURBITS, LEGUME AND FRUITING VEGETABLE, SMALL FRUIT AND BERRIES, ROOT AND TUBER VEGETABLE, ETC. - CONTROL OF MITES**

## General Information

### GENERAL INFORMATION

- Thorough coverage is necessary to provide good disease and mite control. Trilogy prevents fungal attack of plant tissue and contact activity to mite pressure on the plant. Avoid excessive runoff of spray material for maximum control.
- For best results, maintain constant agitation in spray equipment.
- For optimal performance, do not mix with cold water (less than 45°F).
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted crops prior to root establishment.
- Do not apply to known sensitive plant species, such as impatiens flowers, fuchsia flowers, hibiscus flowers, some rose flowers, ornamental olive trees, or some carnation varieties without prior testing.
- When used in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale.
- As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use with care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.

### AGRICULTURAL USE DIRECTIONS

Trilogy is a broad spectrum fungicide of certain diseases and controls mites in citrus, deciduous fruits and nuts, vegetable crops, cereal grains, and other miscellaneous crops.

- Do not use Trilogy after bloom on table grapes or following bunch closure on wine

grapes. Do not use Trilogy after pit hardening on stone fruit.

- Use care when applying near streams, ponds, lakes, or other bodies of water. Do not apply Trilogy when weather conditions favor drift or when the likelihood of runoff is high.

- Apply Trilogy by conventional, low volume, and aerial application equipment.

#### Limitations, Restrictions, and Exceptions

#### CROPS

- Includes cucurbits; legume and fruiting vegetables; small fruit and berries; root and tuber vegetables; bulb, cole and leafy vegetables; and herbs and spices.

- Use the dilution tables to calculate dilution rates (please refer to the label).

- Apply at a rate of 1.0-2.0% Trilogy. For best results, apply Trilogy in sufficient amounts of water to achieve complete coverage. Reapply Trilogy every 7-21 days until pressure is over.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field rates 0](#)

•

#### Restricted Entry Interval

4 hours

#### Timings

[Before mites or eggs are present in large numbers.](#)