

VEGETABLES - STRAWBERRIES, RASPBERRIES, BLUEBERRIES

General Information

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or for other commercial use, or for commercial seed

production or for research purposes.

Do not apply this product in enclosed areas.

Do not use food utensils to mix or apply this product.

Keep people and pets off treated areas until the spray has dried.

SHAKE WELL BEFORE USING

THREE STEPS TO MIX AND APPLY:

1. For lawns and vegetables, measure the area to be sprayed. Calculate square feet by

multiplying length by width. For all other applications, apply as a thorough cover spray.

2. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.

3. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after

each use. Make applications when insects first appear. Repeat applications as directed.

Use Dilution rate as indicated:

1 fl.oz. = 6 tsp. or 2 Tbsp.

8 fl.oz. = $\frac{1}{2}$ pt. or 1 cup

Limitations, Restrictions, and Exceptions

VEGETABLES

Prepare a spray solution at the dilution rates listed for each crop. Apply the spray solution at the rate of 1 gallon to cover 440 square feet surface area.

How to Use

Use 1/2 fl. oz. in 1 gal. of water. Allow 14 days between last application and harvest.

Method

[Foliar application](#)

Pre-Harvest Interval

14 days

Rates

[field rates 0](#)

-

Timings

[N.A.](#)