

CITRUS: (FLORIDA AND TEXAS) - (LOOSENING SOOTY MOLD)

Limitations, Restrictions, and Exceptions

CITRUS: (FLORIDA AND TEXAS) - (LOOSENING SOOTY MOLD)

Apply as a dilute spray.

Comments: Use higher rate when disease pressure is heavy.

Comments: Tank mix with sufficient copper fungicide to provide 4 lbs. per acre of metallic copper.

Dilute spray: Apply the spray oil application rate in up to 4500 gallons of water per acre, or 30 to 35 gallons per tree, to over 45 gallons per tree for large trees. Best results can be obtained when the oil is added to a partially filled spray tank under proper agitation and all foliage is thoroughly covered. Precautions in the use of oil include: Do not apply oil spray when trees are wilting. Do not apply oil and sulfur within 3 weeks of each other in Florida or Texas. Oil spray applied after October 1 may increase susceptibility of trees to cold damage and may reduce the fruit crop the following year. Oil sprays applied in the fall may inhibit solids formation in the juice and retard coloring of fruit and should not be applied within 60 days of anticipated harvest. Sensitive foliage may be injured. Florida growers consult the Florida Citrus Spray and Dust Schedule, or your Agricultural Experiment Station or State Extension Service Specialist for complete details on the spraying program best suited to your locale. Texas growers consult with your Agricultural Experiment Station or State Extension Specialist for complete details on the spraying program suited to your locale.

Method

[Spray](#)

Restricted Entry Interval

4 hours

Timings

[N.A.](#)