

## **APPLICATION - HOPPER BOX TREATMENT**

### General Information

#### ADVERSE CONDITIONS and FIRST-YEAR LAND:

ABM recommends in-furrow use or a double (2X) inoculant rate when planting legumes under the following soil conditions: 1) Soil that has not hosted the specific legume for more than 3 years previously; 2) soil pH of less than 5.8 or greater than 8.5; 3) organic matter less than 1%; 4) drought or flooding; 5) soil temperatures exceeding 80 degrees F; 6) eroded soil; or 7) use of soil treatments and chemicals injurious to soil bacteria and inoculants (see caution below).

#### CAUTION!

- America's Best Inoculant is a living organism.
- All opened containers should be used with 24 hours or less.
- Some seed applied chemicals may be harmful to inoculant bacteria. Check compatibility before use. It is important to protect inoculated seed from excessive heat and rapid drying.

#### Limitations, Restrictions, and Exceptions

#### APPLICATION - HOPPER BOX TREATMENT:

Open box and remove cardboard insert and measuring cup. Pull bag cap through circular opening and close lid. Measure and apply the appropriate amount of America's Best Inoculant for the seed to be treated. Apply uniformly over the top of the seed and stir until all seeds are coated. For optimal performance, plant seed immediately after inoculant application.

#### Method

#### [Seed Treatment](#)

#### Rates

#### [field rates 0](#)

- 

#### Timings

N. A.