

## **DIRECTIONS FOR IN-FURROW USE (7 ROW WIDTH)**

### General Information

#### ADVERSE CONDITIONS and FIRST-YEAR LAND:

ABM recommends in-furrow use or a double (2X) inoculant rate when planting legumes under the following soil conditions: 1) Soil that has not hosted the specific legume for more than 3 years previously; 2) soil pH of less than 5.8 or greater than 8.5; 3) organic matter less than 1%; 4) drought or flooding; 5) soil temperatures exceeding 80 degrees F; 6) eroded soil; or 7) use of soil treatments and chemicals injurious to soil bacteria and inoculants (see caution below).

#### CAUTION!

- America's Best Inoculant is a living organism.
- All opened containers should be used with 24 hours or less.
- Some seed applied chemicals may be harmful to inoculant bacteria. Check compatibility before use. It is important to protect inoculated seed from excessive heat and rapid drying.

#### Limitations, Restrictions, and Exceptions

#### DIRECTIONS FOR IN-FURROW USE:

Apply America's Best Pea & Lentil Inoculant at a rate of 0.5 fl. oz. per 1,000 ft. of row. Shake container well before pouring into mix tank. Dilute with clean, NON-CHLORINATED WATER and use only spray tanks and pumps that are clean and free from chemical contamination. Apply in furrow at a rate of 5 to 20 gallons diluted material per acre. Calibrate pump and spray equipment for the correct delivery rate.

- Unit treats: 2.8 acres

#### Method

#### [In Furrow](#)

#### Rates

## field\_rates 0

- 

Timings

N. A.