

OVERALL LAWN APPLICATION FOR GRASS AND DICHONDRA LAWNS - (BILLBUGS, GRUBS, ETC.)

General Information

LAWNS

Three Easy Steps to use this Product

Step #1: Determine size of the job, measure the area to be treated.

Step #2: Use spreader setting chart in the label to determine appropriate setting for your spreader.

Step # 3: Treat as directed in Rate chart in the label. Thorough coverage is important. Water in granules where indicated.

Note: The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting, if necessary, in order, in order to assure correct rate of application.

Limitations, Restrictions, and Exceptions

OVERALL LAWN APPLICATION FOR GRASS AND DICHONDRA LAWNS - (BILLBUGS, GRUBS, ETC.)

How to Use

Water grass thoroughly after applicaiton. For Billibug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For Grub control, except Dung Beettle, treat any time between late July and earl October,

For Dung Beetle Grub control, apply in spring or early summer when damage is first noticed. Wate grass thouroughly with 1/4 to 1/2 inch of water application. Treat again in late summer or early fall if a second generation occurs.

Method

[Broadcast](#)

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Timings

[N.A.](#)

[N.A.](#)

[N.A.](#)

[N.A.](#)