

SEED TREATMENT

General Information

HDH Peroxy works best when diluted with water containing low levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. HDH Peroxy will readily mix with clean, neutral water and does not require agitation. Do not combine with any other pesticide or fertilizer, HDH Peroxy is formulated with a minimal amount of surfactant for plants having waxy or hairy surfaces. Additional surfactant may be added, if needed. The use of additional surfactant is acceptable. HDH Peroxy works by surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted.

HDH Peroxy does not produce any visible residue, distinct odor or deleterious effects to plants or to postharvest commodities when used in accordance with label directions. Do not use at higher than recommended dilution rates as leaf burn may result. Do not apply this product through any irrigation system unless directed by the label; refer to Chemigation Directions for Use.

Refer the label of agricultural crops.

Limitations, Restrictions, and Exceptions

Seed Treatment -

Use HDH Peroxy for the control of damping-off, root disease and stem rot disease caused by Pythium, Phytophthora, Rhizoctonia, Fusarium or Thielaviopsis, on seeds of seed sprout crops such as mung bean, red clover, soybeans and alfalfa, and on crops grown exclusively for seed for planting.

- 1) Mix 64 fl. oz. HDH Peroxy per 50 gallons of water.
- 2) Immerse seeds and let soak for two minutes; remove and allow to drain. Do not rinse.

Do not use treated seed for food or feed purposes or process for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatments

at planting. Do not store excess treated seeds beyond planting time.

Seed treatments on agricultural establishments in hopper-box, planter-box, or other seed treatment application or immediately before planting is within the scope of WPS, while commercial treatment of seeds is not within the scope.

Method

[Seed Treatment](#)

[Soak](#)

Rates

[field_rates 0](#)

•

Timings

[N.A.](#)