

POTATOES, DIRECT INJECTION

General Information

HDH Peroxy works best when diluted with water containing low levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. HDH Peroxy will readily mix with clean, neutral water and does not require agitation. Do not combine with any other pesticide or fertilizer, HDH Peroxy is formulated with a minimal amount of surfactant for plants having waxy or hairy surfaces. Additional surfactant may be added, if needed. The use of additional surfactant is acceptable. HDH Peroxy works by surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted.

HDH Peroxy does not produce any visible residue, distinct odor or deleterious effects to plants or to postharvest commodities when used in accordance with label directions. Do not use at higher than recommended dilution rates as leaf burn may result. Do not apply this product through any irrigation system unless directed by the label; refer to Chemigation Directions for Use.

Refer the label of agricultural crops.

Limitations, Restrictions, and Exceptions

POTATOES NOTE: Do not treat seed potatoes in storage if sprouting has begun ("Peeps").

Potatoes, direct injection into humidification water for postharvest potatoes in storage. (Fusarium Tuber Rot, Bacterial Soft Rot, Silver Scurf, Early Blight, Late Blight): Inject concentrate into makeup water used in humidification of postharvest potatoes in storage at the rate of 1:100 to 1:300. NOTE: Do not treat seed potatoes if sprouting has begun.

Method

[Inject](#)

Timings

[N.A.](#)