

POTATOES, TREATMENT OF RINSES FOR POSTHARVEST

General Information

HDH Peroxy works best when diluted with water containing low levels of organic or inorganic materials, and with water having a neutral pH. Thoroughly rinse out tank with water before mixing concentrate. HDH Peroxy will readily mix with clean, neutral water and does not require agitation. Do not combine with any other pesticide or fertilizer, HDH Peroxy is formulated with a minimal amount of surfactant for plants having waxy or hairy surfaces. Additional surfactant may be added, if needed. The use of additional surfactant is acceptable. HDH Peroxy works by surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted.

HDH Peroxy does not produce any visible residue, distinct odor or deleterious effects to plants or to postharvest commodities when used in accordance with label directions. Do not use at higher than recommended dilution rates as leaf burn may result. Do not apply this product through any irrigation system unless directed by the label; refer to Chemigation Directions for Use.

Refer the label of agricultural crops.

Limitations, Restrictions, and Exceptions

POTATOES NOTE: Do not treat seed potatoes in storage if sprouting has begun ("Peeps").

Potatoes, treatment of rinses for postharvest potatoes; prior to, during, or after storage. (Odor-causing and/or slime forming bacteria): Inject concentrate into process water, at the rate of 1:1,000 to 1:5000, used in potato rinses and associated tanks, flumes, and lines.

Method

[Inject](#)

Timings

[N.A.](#)