

FIELD AND ROW CROPS: ONION (DRY BULB), GARLIC - BOTRYTIS LEAF BLIGHT OR BLAST, PURPLE BLOTCH (LOW DISEASE HAZARD & PRIOR TO INFECTION)

General Information

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates and wildlife. DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT contaminate water when disposing of equipment washwater or rinsate.

Chlorothalonil can contaminate surface water through spray drift. DO NOT apply when weather conditions favor drift from treated areas. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

Chlorothalonil degradates are known to leach through soil into ground water under certain conditions as a result of label use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

General Precautions and Restrictions

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, or pets, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

Limitations, Restrictions, and Exceptions

APPLICATION DIRECTIONS

CHLOROTHALONIL is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard.

Frequency: 10 days

Method

[Spray](#)

Pre-Harvest Interval

7 days

Rates

[field_rates 0](#)

[field_rates 1](#)

•

Restricted Entry Interval

12 hours

Timings

[Low disease hazard and prior to infection.](#)