

FRUITS AND NUT CROPS: APPLES (WESTERN U.S.) - PRIMARY SCAB

General Information

APPLICATION INFORMATION:

DO NOT apply this product with aerial equipment.

Applications can be made by ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. If necessary, add a few drops of oil to avoid foaming. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read \"COMPATIBILITY AND PLANT SAFETY\" section above.) Application of this material through any type of irrigation equipment is prohibited.

Note: There are 5 level tablespoons or 15 level teaspoons per ounce of this product.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

Do not apply to turfgrass in residential sites, including homes, apartment buildings, daycare centers, schools, playgrounds, parks, recreational areas, sports fields or other residential areas. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control, but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favor disease(s). Apply the low rate and/or spray at longer intervals when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your state Agricultural Extension Service to learn of these conditions.

IMPORTANT: Read label carefully. Although most of the directions on the label may be followed nationwide, a few are limited to either the Eastern or Western U.S. Follow those directions for your growing area where specified.

Limitations, Restrictions, and Exceptions

FRUITS AND NUT CROPS: APPLES (WESTERN U.S.) - PRIMARY SCAB

SPECIFIC DIRECTIONS: Apply in 20 to 400 gals. of water per acre using ground equipment. To reduce potential for tolerance development to other fungicides having a similar spectrum, the lower rate may be used in tank mixtures.

Note: Do not apply more than 64 lbs. of this product per acre from prebloom through postharvest. May be applied up to day of harvest.

Note: Secondary Scab - In mid-Summer cover sprays, the dosage may be reduced to 4 lbs. per acre.

Method

[Broadcast/Foliar Ground](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[N.A.](#)