

POST-HARVEST FRUIT APPLICATION

General Information

APPLICATION INFORMATION:

DO NOT apply this product with aerial equipment.

Applications can be made by ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. If necessary, add a few drops of oil to avoid foaming. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read \"COMPATIBILITY AND PLANT SAFETY\" section above.) Application of this material through any type of irrigation equipment is prohibited.

Note: There are 5 level tablespoons or 15 level teaspoons per ounce of this product.

GENERAL USE PRECAUTIONS AND RESTRICTIONS

Do not apply to turfgrass in residential sites, including homes, apartment buildings, daycare centers, schools, playgrounds, parks, recreational areas, sports fields or other residential areas. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.

Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control, but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favor disease(s). Apply the low rate and/or spray at longer intervals when climatic conditions least favor disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your state Agricultural Extension Service to learn of these conditions.

IMPORTANT: Read label carefully. Although most of the directions on the label may be followed nationwide, a few are limited to either the Eastern or Western U.S.

Follow those directions for your growing area where specified.

Limitations, Restrictions, and Exceptions

POST-HARVEST FRUIT APPLICATION

For use in mechanical fruit-dip operations only. Hand dipping of fruit is prohibited.

For control of various molds and storage rots (*Botrytis*, *Gloeosporium*, *Rhizopus*), use as a postharvest dip or spray wash on the following fruits: Apples, Cherries, Pears — Use 2.5 pounds of this product per 100 gallons of water. Apply as a spray or in a dip tank. When used as a dip, recharge wash solution periodically when tank volume is reduced by 25%. Bring water back to volume and add 2.5 pounds of this product for each 100 gallons added. At end of every 8 hour shift, empty tank, flush and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

Do not contact or allow others to contact the treated fruit until the sprays have dried.

DISPOSAL OF LEFTOVER POSTHARVEST TREATMENT MIXTURE:

Leftover dip or spray mixtures containing captan may be used as a foliar spray for Apples and Cherries, (but not Pears) as treated by the dip or spray mixture or to registered Turf and Ornamental sites. Observe all restrictions such as maximum pounds applied per application and season. When calculating application rates, if analytical services are not available to determine the exact quantity of captan remaining in the mixture, assume that the tank still contains 2.5 pounds of this product per 100 gallons of water. If the dip or spray mixture contains other pesticide(s) in addition to captan, refer to the product label(s) for information regarding disposal.

Captan wastes are acutely hazardous to the eyes. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. Contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance concerning the disposal of spent or excess dip tank mixtures.

Method

[Dip](#)

[Spray](#)

Rates

[field_rates 0](#)

•

Timings

[Post-harvest](#)