

## **FOLIAR APPLICATIONS - DILUTE APPLICATION**

### General Information

#### GENERAL INFORMATION

DESERT MICRO BLEND is especially recommended to prevent and correct copper, iron, manganese and zinc deficiencies in desert-grown field, row, vegetable, fruit, tree, and vine. DESERT MICRO BLEND alone, with pesticides, or with other fertilizers, may be applied directly to the soil or to the foliage of growing crops. DESERT MICRO BLEND may be applied via ground sprayers (high-volume or lowvolume), aircraft or fertigation systems.

#### DIRECTIONS FOR USE

Soil applications of DESERT MICRO BLEND are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results.

Actively growing plants with young leaves absorb foliarly applied DESERT MICRO BLEND most efficiently. Immediate post-irrigation or post-rainfall application give the best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Bi Urea per 100 gallons of spray solution) will enhance foliar uptake of DESERT MICRO BLEND.

#### Limitations, Restrictions, and Exceptions

#### FOLIAR APPLICATIONS FOR ALL CROPS:

Applications should be separated by a minimum of 7-14 days. AVOID foliar applications of this product when plants are under moisture stress. AVOID foliar applications to deciduous tree crops during bloom.

Dilute Applications: For rates higher than 2 quarts per acre, split applications of 2 quarts or less are preferred to minimize phytotoxicity.

#### Method

[Foliar spray](#)

#### Rates

[field\\_rates 0](#)

- 

Timings

[N.A.](#)