

## **SOIL APPLICATION - LIGHT DEFICIENCY**

### General Information

### General Information

FERTI-man 6% is recommended for field, row, vegetable, fruit, tree, vine, ornamental, and turf crops.

FERTI -man 6%, alone or with pesticides or fertilizers, may be applied directly to soil or to the foliage of growing crops.

FERTI-man 6% may be applied via ground sprayers (high-volume or low-volume), aircraft or drip irrigation systems.

### General Directions for use

Consult your local Fertizona representative or other agricultural specialist for specific crop recommendations.

Soil application of FERTI-man 6% is recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly applied FERTI-man 6% most efficiently. Immediate post-irrigation or post-rainfall applications give best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Biuret Urea per 100 gals. of spray solution) will enhance foliar uptake of FERTI-man 6%.

Avoid foliar applications of FERTI -man 6% when plants are under moisture stress.

Avoid application to deciduous tree crops during bloom.

### DILUTE APPLICATIONS

Apply the suggested dose of Ferti-Man 6% in enough water for good coverage.

### DRIP IRRIGATION APPLICATIONS

Apply the recommended amount of Ferti-Man 6% in split applications 1-2 weeks apart throughout the growing season.

## Limitations, Restrictions, and Exceptions

- The lower rates are for field, row, and vegetable crops. The higher rates are for fruit, tree, vine, ornamental, and turf crops.

### Method

[Soil application](#)

### Rates

[field rates 0](#)

- 

### Timings

[N.A.](#)