

FOLIAR APPLICATION

General Information

General Information

FERTI-man 6% is recommended for field, row, vegetable, fruit, tree, vine, ornamental, and turf crops.

FERTI -man 6%, alone or with pesticides or fertilizers, may be applied directly to soil or to the foliage of growing crops.

FERTI-man 6% may be applied via ground sprayers (high-volume or low-volume), aircraft or drip irrigation systems.

General Directions for use

Consult your local Fertizona representative or other agricultural specialist for specific crop recommendations.

Soil application of FERTI-man 6% is recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Actively growing plants with young leaves absorb foliarly applied FERTI-man 6% most efficiently. Immediate post-irrigation or post-rainfall applications give best foliar uptake. Addition of urea (for example, 5 lbs. Lo-Biuret Urea per 100 gals. of spray solution) will enhance foliar uptake of FERTI-man 6%.

Avoid foliar applications of FERTI -man 6% when plants are under moisture stress.

Avoid application to deciduous tree crops during bloom.

DILUTE APPLICATIONS

Apply the suggested dose of Ferti-Man 6% in enough water for good coverage.

DRIP IRRIGATION APPLICATIONS

Apply the recommended amount of Ferti-Man 6% in split applications 1-2 weeks apart throughout the growing season.

Limitations, Restrictions, and Exceptions

FOLIAR APPLICATIONS

For rates (above) higher than 2 qts. per acre, split applications of 2 qts. or less each are preferable. Applications should be separated by a minimum of 10-14 days.

Avoid foliar applications of this product when plants are under moisture stress.

Avoid application to deciduous tree crops during bloom.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field_rates 0](#)

-

Timings

[N.A.](#)