

TURFGRASSES - WARM SEASON GRASSES - CUDWEED ETC. - ALL TURF USES

General Information

USE INFORMATION

PENDIGUARD TURF 0.75% is a combination preemergent herbicide with fertilizer formulated to provide control of most annual grass weeds and certain broadleaf weeds found in Turfgrass grown in golf courses, lawns in residential, commercial establishments, or grounds around military or other institutions, sod farms, parks, airports, school or cemeteries, prairiegrass areas, athletic fields or recreational areas, roadsides, pipe yards, industrial, unimproved Turf areas, and other sites where Turf may be maintained. This product is labeled for use on the following Turfgrass species: Kentucky bluegrass, Perennial ryegrass, Fine fescue, Tall fescue, Bentgrass (except putting greens and tees), Poa annua (except putting greens and tees), Bermudagrass, St. Augustinegrass, Bahiagrass, Centipedegrass, Zoysiagrass, and blends of these grasses.

This product controls weeds as their seeds germinate, but will not control established weeds. Use a postemergent herbicide, such as 2,4-D or MSMA, to control established weeds. Consult all herbicide label directions for proper use and precautions.

Applied according to label directions and under normal growing conditions, this product will not cause Turf injury. Over application can result in Turf stand loss, Turf injury, or soil residues. Uneven application can decrease weed control or cause Turf injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, soil salt concentration, or drought can weaken Turfgrass and increase the possibility of Turf damage from this product.

Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

This product can be used around established Ornamentals and in established

Wildflower plantings. This product may be used for hardwood and conifer regeneration on conservation reserve program land.

APPLICATION

Apply this product uniformly at the listed rate using a properly calibrated spreader (see table below for approximate settings). Avoid overlaps that will increase rates above those listed. The efficacy of this product will be improved if the application is followed by one-half inch of rainfall or its equivalent in sprinkler irrigation. This will remove the product from the Turfgrass foliage and establish a preemergent herbicide barrier in the soil. If this product is not activated by rainfall or irrigation within 30 days of application, erratic weed control may result. Avoid contact of this product with porous surfaces such as driveways, stone, or wood. Removing particles will help prevent staining.

Refer the Maximum Application Rate By Use Sites.

Limitations, Restrictions, and Exceptions

COMMENTS:

- Apply a repeat application of 200 to 267 pounds per acre after 5 to 8 weeks for extended Poa annua control.

Method

[Broadcast](#)

Rates

[field rates 0](#)

-

Restricted Entry Interval

24 hours

Timings

[In late summer or early fall prior to weed germination.](#)