

## **AIR OR GROUND**

### General Information

### DIRECTIONS FOR USE

Consult your local Fertizona representative or other agricultural specialist for specific crop recommendations.

SUPR-SPRED is a non-ionic surfactant containing an anti-foaming agent. It improves wetting of foliage and enhances the effectiveness of spray applications. It will not cause excessive foaming in the spray tank nor corrode equipment. SUPR-SPRED may be used on agricultural, aquatic, forestry, turf and ornamental, industrial, structural and non-cropland sites.

Read and follow the precautions, restrictions and recommendations on the labels of pesticides used with SUPR-SPRED. Use according to the most restrictive label directions for each product in any tank mix.

**COMPATIBILITY:** SUPR-SPRED is compatible with most pesticides. However, if the desired combination has not been previously used, a compatibility test is recommended.

**MIXING:** Shake well before using. Fill spray tank  $\frac{1}{2}$  full of water and begin agitation. Add the recommended amount of SUPR-SPRED. Add products as directed by label or in the following sequence and continue filling tank: (1) Dry flowables or water dispersible granules, (2) Wettable powders, (3) Flowables, (4) Solutions, (5) Emulsifiable concentrates. Continue agitation until spray solution is completely mixed. Continuous agitation of finished spray is recommended. If spray solution has been allowed to stand, thoroughly agitate and remix before application.

### Limitations, Restrictions, and Exceptions

**RECOMMENDATIONS:** Specific use rates will vary with conditions of application such as water hardness, application method, equipment, spray droplet size, condition of foliage, etc. Also, higher rates than those below may be used if recommended by pesticide labeling. Follow pesticide label directions. However, do not add this

product at a rate which exceeds 5% of the finished spray volume.

- AIR or GROUND: Use 0.5-4 pints per 100 gallons of finished spray solution. For improved efficacy, use at least 0.5 pint per acre when finished spray volume is less than 20 gallons per acre. For backpack or hand held sprayers, use 0.5 fl.oz. per gallon of finished spray solution. Exercise caution with selective herbicides where selectivity depends upon wetting the weeds but not the crop.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Rates

[field\\_rates 0](#)

•

Timings

[N. A.](#)