

FOR CONTROL OF HOUSE MICE

General Information

USE RESTRICTIONS

This product can only be used to control Norway rats, roof rats, and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Burrow baiting with Rodenthor Block Bait is prohibited.

SELECTION OF TREATMENT AREAS

After removing as much food, water and harborage as possible, determine dry areas where rats and mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats or house mice have been seen. Protect bait from rain and snow.

APPLICATION DIRECTIONS

Each RODENTHOR WAX BLOCK weighs approximately 1.0 oz.

Follow-up: While wearing gloves, replace contaminated or spoiled bait immediately. Collect and properly dispose of dead animals and leftover bait and bait bags, as well as any bait that has become accessible to non-target animals. To discourage re-infestation, limit sources of rodent food, water, and harborage as much as possible.

If re-infestation does occur, repeat treatment. For continuous sources of infestation, set up permanent bait stations and replenish them as needed.

Limitations, Restrictions, and Exceptions

FOR CONTROL OF HOUSE MICE

Maintain a constant supply of bait blocks for 15 days or until signs of mouse activity cease.

Method

[Bait](#)

Rates

[field_rates 0](#)

-

Timings

[N. A.](#)