

BROADCAST APPLICATION FOR ORNAMENTAL PESTS

General Information

DIRECTIONS FOR USE

Do not apply more than 80 pounds (0.4 pounds of active ingredient) of TURFTHOR 0.5 G per acre per year for any type(s) or combination(s) of applications.

Do not apply in greenhouses or nurseries.

Not for use on sod farm turf or grass grown for commercial seed production.

Not for use on bearing fruit or nut trees.

RESTRICTIONS

Do not graze treated areas or use clippings from treated areas for feed or forage.

Do not allow runoff or puddling of irrigation water applied following application.

Limitations, Restrictions, and Exceptions

BROADCAST APPLICATION FOR ORNAMENTAL PESTS

- TURFTHOR 0.5 G, broadcast on the ground, controls a wide range of insects on trees (including non-bearing fruit), shrubs, evergreens, flowers, foliage plants, ground covers and interior plantscapes. Non-bearing trees are perennial plants that will not produce a harvestable agricultural commodity within the next 12 months. TURFTHOR 0.5 G is not for use on plants being grown for sale, fruit, nut or commercial seed production or for research purposes.

- TURFTHOR 0.5 G is a systemic insecticide meaning it will be translocated by the plant's vascular system from the roots up into the body of the plant. This means optimum effectiveness of TURFTHOR 2.5 G is realized when TURFTHOR 2.5 G is applied on the ground above the root zone of the plant to be protected. Watering after application can speed movement of the active ingredient into the plant's vascular system. Combining TURFTHOR 0.5 G with a nitrogen containing fertilizer may further accelerate or otherwise enhance the uptake of the active ingredient

into the plant.

Translocation of soil directed applications made to woody stemmed plants can be delayed by up to 60 days. Optimum levels of control are realized when applications are performed sufficiently prior to anticipated pests infestations.

Application Sites

- For use on ornamental plants including but not limited to trees (including non-bearing fruit), shrubs, evergreens, flowers, ground covers, bedding plants and foliage plants. Permitted sites include landscapes, ornamental gardens, lawns, grounds, golf courses and interior plantscapes at and/or around residences including multi-unit, commercial, office and shopping buildings and complexes, recreational areas, playgrounds, athletic fields, golf courses, cemeteries, airports and parks, public and private wooded and forested areas.

Application Precautions and Preparations

- Avoid application to areas in which penetration to the root zone is unlikely to occur such as areas that are waterlogged, saturated or frozen.

- TURFTHOR 0.5 G can be mixed with other granular insecticides, miticides, fungicides, herbicides and fertilizers. Follow the label directions of all the products mixed, making sure not to exceed the labeled application rate of any individual product in the mixture.

- If necessary, consult resources in horticulture in your area (such as your Cooperative Extension Service) to determine appropriate application timing.

Post Application Watering

- Optimum control is achieved if irrigation or rainfall occurs within 24 hours after application.

Application Equipment and Methods

- Apply TURFTHOR 0.5 G to the ground around ornamentals using properly calibrated granular application equipment designed to distribute the active ingredient to the target area(s) at the indicated rate(s). Drop-type and rotary-type granular spreaders can be used to apply Turfthor 0.5 G.

Method

[Broadcast](#)

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

[Prior to anticipated pests infestations.](#)