

AVOCADOS AND CITRUS FRUIT

General Information

General Information

CORE Zinc 10% is an effective, readily available source of liquid Zinc for both soil and foliar applications. It may be applied in water sprays, with pesticides, or with most liquid fertilizers.

Directions for Use

Proper timing, rate, and placement of CORE Zinc 10% is important for desired results and highly dependent on stage of crop growth, soil fertility levels, and environmental conditions. Over application of CORE Zinc 10% may be phytotoxic or cause undesirable effect on foliage. For foliar applications, apply with enough water to thoroughly cover all foliage. Early morning or late evening applications generally give the best results. For aerial application, use at least one gallon of water for each pint of product applied. For soil applications, apply with sufficient carrier to give complete coverage. For optimum performance, this can be accomplished by irrigation, rainfall, or tillage. Do not allow CORE Zinc 10% to come in direct contact with seed at any time.

- The product is recommended to be applied early in the growing season.

Limitations, Restrictions, and Exceptions

Apply 1 to 2 gallons per acre per application. Apply first application at pre-bloom, then subsequent applications at mid crop and 3 to 4 weeks prior to harvest, or as needed to meet nutritional requirements.

Method

[Foliar spray](#)

[Soil application](#)

Rates

[field rates 0](#)

-

Timings

Pre-bloom.