

# STONE FRUIT

## General Information

## General Information

CORE Zinc 12% is an effective, readily available source of liquid Zinc for both soil and foliar applications. It may be applied in water sprays, with pesticides, or with most liquid fertilizers.

## Directions for Use

Proper timing, rate, and placement of CORE Zinc 12% is important for desired results and highly dependent on stage of crop growth, soil fertility levels, and environmental conditions. Over application of CORE Zinc 12% may be phytotoxic or cause undesirable effect on foliage. For foliar applications, apply with enough water to thoroughly cover all foliage. Early morning or late evening applications generally give the best results. For aerial application, use at least one gallon of water for each pint of product applied. For soil applications, apply with sufficient carrier to give complete coverage. For optimum performance, this can be accomplished by irrigation, rainfall, or tillage. Do not allow CORE Zinc 12% to come in direct contact with seed at any time.

- The product is recommended to be applied early in the growing season.

## Limitations, Restrictions, and Exceptions

Apply 4 to 8 gallons per acre per application. Apply first application at green tip or pink bud. Apply subsequent applications at fruit set or jacket fall, fruit thinning and 2 to 3 weeks prior to harvest, or as needed to meet nutritional requirements.

## Method

[Foliar spray](#)

[Soil application](#)

[Foliar spray](#)

[Soil application](#)

## Rates

## field\_rates 0

- 

Timings

At green tip or pink bud.

At fruit set or jacket fall, fruit thinning and 2 to 3 weeks prior to harvest.