

## **APPLES, PEACHES, PEARS, ETC.**

### General Information

### SUGGESTED DIRECTIONS

General use: Apply 5 to 10 lbs. per acre as a foliar application to supplement the normal soil applied fertility program. Can be applied by aircraft at rates equivalent to 1 lb. in 1 gal. of water, per acre. For back-pack sprayers use at the rate of 6 grams per liter or 4 teaspoonfuls per gallon.

- 5 LBS. PER ACRE = 5 KG PER HECTARE CAN BE USED WITH NU-FILM-P OR NU-FILM-17

COMPATIBILITY: NUTRI-LEAF is compatible with most pesticides commonly used on vegetable, fruit and agronomic crops. CAUTION: Do not use with Dormant Oil, Lime Sulfur, Spray Lime, or highly alkaline spray materials. Add NUTRI-LEAF to the spray tank or pre-mix tank when it is about half full. Agitation in the tank is preferred to increase the speed of solubility. Where extremely cold water is used, allow a few extra minutes for the product to completely dissolve before spraying.

### Limitations, Restrictions, and Exceptions

APPLES, PEACHES, PEARS, PLUMS, NECTARINES, APRICOTS, CHERRIES, CITRUS FRUITS, FIGS, AVOCADOS, MANGOES, PAPAYA, KIWI, FILBERTS, CHESTNUTS, MACADAMIA NUTS, WALNUTS, PECANS AND ALMONDS: Use at the rate of 2 lbs. per 100 gals. of water when applying in dilute sprays or 5 to 10 lbs. per acre when applying with concentrate sprayers. Apply early in the growing season and whenever supplemental feeding is necessary, to improve tree vigor, fruit or nut set or during adverse weather conditions. Can be applied in combination with almost all insecticide and fungicide programs. For most tree crops, 3 to 5 applications are desirable through the growing season. CAUTION: On tree fruits, where fruit color and maturity is delayed by nitrogen application, do not use NUTRILEAF in late season sprays.

### Method

#### [Spray](#)

Rates

[field\\_rates 0](#)

- 

Timings

[Apply early in the growing season and whenever supplemental feeding is necessary.](#)