

STRAWBERRIES

General Information

SUGGESTED DIRECTIONS

General use: Apply 5 to 10 lbs. per acre as a foliar application to supplement the normal soil applied fertility program. Can be applied by aircraft at rates equivalent to 1 lb. in 1 gal. of water, per acre. For back-pack sprayers use at the rate of 6 grams per liter or 4 teaspoonfuls per gallon.

- 5 LBS. PER ACRE = 5 KG PER HECTARE CAN BE USED WITH NU-FILM-P OR NU-FILM-17

COMPATIBILITY: NUTRI-LEAF is compatible with most pesticides commonly used on vegetable, fruit and agronomic crops. CAUTION: Do not use with Dormant Oil, Lime Sulfur, Spray Lime, or highly alkaline spray materials. Add NUTRI-LEAF to the spray tank or pre-mix tank when it is about half full. Agitation in the tank is preferred to increase the speed of solubility. Where extremely cold water is used, allow a few extra minutes for the product to completely dissolve before spraying.

Limitations, Restrictions, and Exceptions

STRAWBERRIES: Use at the rate of 5 to 10 lbs. per acre when the first fruit buds show in the crown. Three or more additional applications are desirable at 7 to 10 day intervals.

Method

[Spray](#)

Rates

[field rates 0](#)

-

Timings

[When the first fruit buds show in the crown.](#)