

COTTON

General Information

SUGGESTED DIRECTIONS

General use: 5 to 10 lbs. per acre. Use this product as a foliar application to supplement the normal soil applied fertility program. Can be applied by aircraft at rates equivalent to 1 lb. in 1 gal. of water, per acre. For back-pack sprayers use at the rate of 6 grams per liter or 4 teaspoonfuls per gallon.

- CAN BE USED WITH NU-FILM-P or NU-FILM-17

COMPATIBILITY: NUTRICHEM 9-15-30 is compatible with most pesticides commonly used on vegetable, fruit and agronomic crops. CAUTION: Do not use with Dormant Oil, Lime Sulfur, Spray Lime, or highly alkaline spray materials. Add NUTRICHEM 9-15-30 to the spray tank or pre-mix tank when it is about half full. Agitation in the tank is preferred to increase the speed of solubility. Where extremely cold water is used, allow a few extra minutes for the product to completely dissolve before spraying.

TRANSPLANTING: Use 5 lbs. per 100 gallons of water. (1 cup of mix per plant for hand setting) It requires about 200 to 300 gallons of water per acre for transplanting, or 10 to 15 lbs. of NUTRICHEM 9-15-30 per acre.

Limitations, Restrictions, and Exceptions

COTTON: To stimulate boll development, maturity, and fiber strength, apply NUTRICHEM 9-15-30 as a foliar spray at the rate of 5 to 10 lbs. per acre. Applications should begin directly after peak bloom, and continue at 7 to 10 day intervals. Apply 3 to 4 applications for best results.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

•

Timings

After peak bloom