

# **APPLES, CRAB APPLES, LOQUATS, PEARS AND QUINCE - APPLE BLACK SPOT AND SCAB**

## General Information

### GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

### Limitations, Restrictions, and Exceptions

### APPLICATION PROGRAM

Total of 10 applications at 10 to 12 day intervals. When conditions are conducive to a black spot outbreak, apply ORGANOCIDE PLANT DOCTOR immediately. Note: After 4 or 5 consecutive applications some yellowing of extension growth may be observed. If yellowing occurs use another fungicide until yellowing of leaves disappears.

Not for use in California except as basal bark spray on apples and crab apples.

### Method

[Foliar spray](#)

### Rates

[field\\_rates 0](#)

- 

### Timings

[First application at open cluster. Last application at fifth cover or fruit at 2" to 2 1/2" diameter.](#)