

AVOCADOS - ROOT ROT - OTHER SITUATIONS

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

Limitations, Restrictions, and Exceptions

APPLICATION PROGRAM

Repeat treatment in February or March. Drill holes 3/16 inch (5 mm) in diameter and 1 inch (25 mm) to 2 inches (50 mm) deep with slight downward angle in trunk. Place syringes in the main trunk of the tree and space evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun, or hydraulic tree injection. Do not prune back trees before injection process as burning of new growth may occur. Do not inject trees in winter months. Do not cut back the canopy of injected trees. Do not add any material, other than water, to ORGANOCIDE PLANT DOCTOR by trunk injection. Do not inject more liquid in a lesser number of syringes than directed.

Method

[Injection](#)

Rates

[field rates 0](#)

-

Timings

[At spring flush maturity.](#)