

GRAPES - DOWNY MILDEW

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

Limitations, Restrictions, and Exceptions

APPLICATION PROGRAM

It is essential that the rate of ORGANOCIDE PLANT DOCTOR be adjusted to the vine-row volume, i.e., the volume of vine foliage per acre. Spray timing is critical. Ensure spray coverage is adequate and that the appropriate rate of ORGANOCIDE PLANT DOCTOR is applied to match vine growth, particularly from mid-season onwards, and especially where grapes are grown on root stock.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

-

Timings

[At times of high disease risk, especially between the time that conditions are conducive to downy mildew infection and the appearance of oil spots.](#)