

LEAFY VEGETABLES

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

Limitations, Restrictions, and Exceptions

LEAFY VEGETABLES

Use ORGANOCIDE PLANT DOCTOR for effective control of downy mildew in leafy vegetables such as, but not limited to, amaranth, arugula, cardoon, celery, chervil, corn salad, endive, fennel, lettuce, parsley, radicchio, rhubarb, spinach, and Swiss chard. Excludes Brassica vegetables.

Use in California limited to lettuce.

APPLICATION PROGRAM

Ensure spray coverage is adequate to wet the whole plant.

During warm, wet conditions repeat application at 7 to 10 day intervals, if needed.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

-

Timings

During warm, wet conditions.