

ORNAMENTAL APPLICATIONS - DOWNY MILDEW

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

Limitations, Restrictions, and Exceptions

ORNAMENTAL APPLICATIONS

Use ORGANOCIDE PLANT DOCTOR for effective control of Bacterial blight, Downy mildew, Phytophthora spp [Root Rot], and Pythium spp [Damping-off] diseases, and Sudden Oak Death of Ornamentals in landscapes and interiorscapes. Apply ORGANOCIDE PLANT DOCTOR to plants such as, but not limited to, Aglaonema, Aphelandra, Arborvitae, Azaleas, Bougainvillea, Boxwood, Cattelya skinneri, Ceanothus, Cotoneaster, Cissus, Diffenbachia, English ivy, Eucalyptus, Ficus, Hibiscus, Japanese andromeda, Japanese Holly, Leather leaf Fern, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Rhododendron, Roses (container, landscape, mini varieties), Schefflera, Sedum, Sempervivum, Syngonium, Spathiphyllum, Taxus media, and Zygocactus.

Make applications before disease development and in conjunction with good cultural management practices. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to plants that are heat or moisture stressed. Do not apply to plants that are in a state of dormancy. When applying to indoor plants do not overspray and use care to apply only to target plants. If meeting these conditions is not possible, remove plants to

an outdoor location for treatment and drying before bringing back indoors.

APPLICATION PROGRAM

Apply spray to thoroughly wet all foliage. Application intervals: 14 to 21 days.
Repeat as required.

Method

[Foliar spray](#)

Rates

[field rates 0](#)

-

Timings

[Before disease development and in conjunction with good cultural management practices.](#)