

CONIFERS - FUSARIUM SUBGLUTINANS - BASAL BARK SPRAY

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

Limitations, Restrictions, and Exceptions

CONIFERS

Apply ORGANOCIDE PLANT DOCTOR in conjunction with good cultural management practices for effective control of root rot (*Phytophthora* spp) in CONIFERS including, but not limited to, Pines, Spruce and Douglas Fir. Use higher rate of application when disease pressure is severe. Do not exceed indicated application rates or apply more frequently than stated on label in order to avoid plant injury. Do not apply to CONIFERS that are moisture or heat stressed.

APPLICATION PROGRAM

- Add 0.5 fl. oz. of Pentra-Bark Bark Penetrating Surfactant.

Apply uniformly to 6-9 feet of trunk circumference. Spray from top down to ground level from either first branch or from as high as possible without exposing applicator to drift. Spray to saturation/ runoff. Various types of application equipment can be used such as hydraulic sprayers, handheld pump-type sprayers, and backpack sprayers

- Use in California limited to injection and/or basal bark spray of pines.

Method

Spray

Rates

field rates 0

•

Timings

Before disease development and in conjunction with good cultural management practices.

Any time active growth occurs.