

TURF

General Information

GENERAL APPLICATION INSTRUCTIONS

Apply ORGANOCIDE PLANT DOCTOR by various application methods, including foliar spray, soil drench, soil incorporation, basal bark application and bare root dip. For foliar sprays, apply ORGANOCIDE PLANT DOCTOR with sufficient water volumes for adequate coverage of foliage, according to plant type and growth stage. To ensure good coverage, spray to wetness, but avoid run-off.

When applying ORGANOCIDE PLANT DOCTOR to plant species for the first time, spray a limited number of plants first and wait for 3-7 days. Then check for leaf burn (phytotoxicity).

Limitations, Restrictions, and Exceptions

TURF

Use ORGANOCIDE PLANT DOCTOR for the effective control of Pythium and damping-off diseases of turf grasses. When conditions favor disease, begin preventive applications and repeat at indicated intervals. Use higher rate of application when disease pressure is severe.

APPLICATION PROGRAM

Apply indicated quantity of product in 1 to 5 gal. of water per 1000 sq. ft. Ensure foliage is thoroughly wet. Application intervals: 14-21 days. Repeat as required. Do not irrigate or mow treated areas until spray has completely dried.

Do not graze animals on treated areas of turf. Do not feed treated turf clippings to poultry or livestock.

Method

[Drench](#)

Rates

[field rates 0](#)

-

Timings

Before disease development and in conjunction with good cultural management practices.