

# **TURFGRASS APPLICATIONS: ANNUAL BLUEGRASS WEEVIL, ASIATIC GARDEN BEETLE, ETC.**

## General Information

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTE: Not For Use in New York State or California

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

Apply only when wind speed is 10 mph or less at the application site.

### Limitations, Restrictions, and Exceptions

#### Application:

Apply Sharda Imidacloprid 70 WDG in sufficient water to provide adequate distribution in the treated area. Use a low pressure setting, and apply a uniform coarse droplet spray, which will eliminate off-target drift. Check equipment calibration periodically to ensure that the product is being distributed properly.

### APPLICATION INSTRUCTIONS

- For best control, apply before egg hatch.
- Chinch bugs (suppression): apply the maximum rate before the first instar nymphs hatch.

- Mole crickets: apply before or during the peak egg hatch period. For adults or large nymphs that are actively tunneling, combine Sharda Imidacloprid 70 WDG with an insecticide with a different mode of action. Follow label instructions for other insecticides when tank mixing.

- To ensure that the active ingredient moves into the turf, irrigation or rainfall must occur within 24 hours of application. Do not mow sufficient irrigation or rainfall has occurred.

## Method

[Spray](#)

## Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

## Timings

[Mole crickets: Before or during the peak egg hatch period.](#)

[Before egg hatch.](#)

[Chinch bugs \(suppression\): Before the first instar nymphs hatch.](#)