

APPLE, CRABAPPLE, ETC.: APHIDS, LEAFHOPPER, ETC.

General Information

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTE: Not For Use in New York State or California

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

Apply only when wind speed is 10 mph or less at the application site.

Limitations, Restrictions, and Exceptions

FOR USE ON POME FRUITS IN AND ON RESIDENTIAL AREAS ONLY

(Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, Quince)

PESTS:

- Aphids (except Woolly apply aphid)
- Leafhoppers (incl. glassy-winged sharpshooter)
- Mealybugs (not for use on this pest on pears in California)
- San Jose Scale (not for use on this pest on pears in California)

RATE:

- 0.5 oz. (3 tbsp. + 1 tsp.) per 100 gallons or 2 oz./acre

Note: The amount of product per acre will vary according to tree size and volume of foliage. The per acre rate is based on a standard 400 gallons dilute solution per acre for large trees.

APPLICATION INSTRUCTIONS:

- Apply at the specified rate as needed once petal fall is complete.
- Do not apply more than 2 ounces per acre in one single application. Do not make more than 5 applications.
- Do not apply more often than every 10 days.

Rosy apply aphid: Apply before the pest begins leafrolling.

First generation leafminers: Apply as soon as petal fall is complete. For best control, apply as early as possible.

Second and later generation leafminers: Apply early in the adult flight to control eggs and early instar larvae. If necessary under conditions of severe pressure or overlapping generations, make a second application. This product will not control late stage larvae.

San Jose Scale: Make application at the crawler stage. Treat each generation.

Leafhoppers, late season control: Apply when most leafhoppers are nymphs.

Mealybug: For best control, good coverage of trunk, scaffolding limbs, or other resting sites is essential.

Method

[N.A.](#)

Pre-Harvest Interval

7 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Timings

San Jose Scale: At the crawler stage.

First generation leafminers: As soon as petal fall is complete.

Second and later generation leafminers: Early in the adult flight to control eggs and early instar larvae.

Leafhoppers, late season control: When most leafhoppers are nymphs.

Once petal fall is complete.

Rosy apply aphid: Before the pest begins leafrolling.