

## **FOAM APPLICATIONS (2.5 OZ)**

### General Information

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTE: Not For Use in New York State or California

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

Apply only when wind speed is 10 mph or less at the application site.

#### Limitations, Restrictions, and Exceptions

#### FOAM APPLICATIONS

In some cases construction practices, soil subsidence and other factors may make it difficult to create a continuous chemical treatment zone using conventional treatment alone. In situations where necessary, supplement conventional application methods by use of foam generating equipment, or similar devices, to create a continuous treated zone. Foam application may be made alone or combined with conventional application methods, provided that the labeled amount of active ingredient per unit area is used.

Foam Application Use Directions: Mix solution of Sharda Imidacloprid 70 WDG with manufacturer's specified volume of foaming agent (see table for foaming directions). Apply a sufficient volume of Sharda Imidacloprid 70 WDG foam alone or in combination with liquid solution to create a continuous treated zone at the specified rate for specific application sites. Use appropriate dispersion tips and application method for site.

Depending on the circumstances, foam applications may be used alone or combined with liquid solution applications. Applications may be made behind

veneers, piers, chimney bases, into rubble foundations, into block voids or structural voids, wall voids, under slabs, stoops, porches, or to the soil in crawl spaces, and other similar voids.

Foam and liquid applications must be consistent with volume and active ingredient instructions in order to ensure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the gallons of Sharda Imidacloprid 70 WDG must be applied as a typical liquid treatment. The remaining 25% or less gallons are delivered to appropriate location application.

NOTE: When foam is used solely to kill subterranean termites above ground locations (such as feeding galleries in wooden framing, or in voids with framed walls), and whenever the target pest is other than subterranean termites (drywood termites, ants, etc.), dilute solutions of imidacloprid may be expanded by foaming without concentrating the imidacloprid solution as previously described for soil applications. Add the manufacturer's specified volume of foaming agent to produce foam of the desired expansion ratio. Use application suitable to the site and pest.

Note: Use Rate - 2.5 oz

1 gal water - 25:1 (Foam Expansion Ratio) or 2.5 gal water - 10:1 (Foam Expansion Ratio) or 5 gal water - 5:1 (Foam Expansion Ratio)

25 gallons/0.05 ai% (Finished Foam)

- Add the manufacturer's specified quantity of foam agent to the Sharda Imidacloprid 70 WDG solution.

Method

[Foam Application](#)

Rates

[field\\_rates 0](#)

•

Timings

[N. A.](#)