

## **HERBS (EXCEPT CHIVES)**

### General Information

Resistance Management: Because of the inherent risks of resistance development to any product, it is strongly advised that Solera Abamectin Agricultural Miticide/Insecticide be used in a sound resistance management program that includes rotation with other products with different modes of action.

### Limitations, Restrictions, and Exceptions

#### HERBS (EXCEPT CHIVES)

#### Application Instructions

Mix with water as indicated below. Apply thoroughly with ground application equipment as a foliar spray and ensure good upper and lower leaf coverage.

- Use 8 fl. oz./A for low to moderate infestations and 16 fl. oz./A for severe infestations.

- Apply when adult flies are first observed and repeat application, as necessary to maintain control. (See Use Restrictions)

Note: Solera Abamectin Agricultural Miticide/Insecticide may be used without a wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Note: Although Solera Abamectin Agricultural Miticide/Insecticide has been tested in combination with a nonionic surfactant for safety to herbs, it is impossible to test on all herb types and varieties under the variety of conditions that may cause crop injury. Therefore, when using Solera Abamectin Agricultural Miticide/Insecticide in combination with a nonionic surfactant, carefully follow the Directions for Use and Precautions on the nonionic surfactant label and in official spray guides.

#### Use Restrictions - Herbs

- Apply with ground application equipment.

- Use a minimum of 30 gals of water/A.
- Wait at least 7 days before repeating application.
- Do not make more than 2 applications per single cutting (harvest).
- Do not make more than 6 applications per cropping season at the 8 fl. oz./A rate or 3 applications per cropping season at the 16 fl. oz./A rate.

#### Method

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

14 days

#### Rates

[field rates 0](#)

- 

Restricted Entry Interval

12 hours

Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4-days.

#### Timings

[when adult flies are first observed](#)