

MINT (PEPPERMINT AND SPEARMINT)

General Information

Resistance Management: Because of the inherent risks of resistance development to any product, it is strongly advised that Solera Abamectin Agricultural Miticide/Insecticide be used in a sound resistance management program that includes rotation with other products with different modes of action.

Limitations, Restrictions, and Exceptions

MINT

Application Instructions

- Mix with water as indicated below. Thorough coverage of the crop canopy is essential for optimum results. Inadequate coverage may result in reduced control.
- Treat when mites first appear.
- For optimum control, a nonionic surfactant or organosilicone-based surfactant should be used at the manufacturer's recommended rate.

Use Restrictions - Mint

- Ground or aerial application permitted; however do not apply with aircraft in New York State.
- Do not apply in less than 5 gals. of water/A with aircraft.

Under conditions such as high pest populations, dense foliage, or adverse application conditions such as high temperatures, use a greater volume of water to insure adequate coverage.

- Wait at least 7 days before repeating application.
- Do not make more than 2 sequential applications and do not apply more than 3 applications per season.
- Do not allow livestock to graze or feed treated foliage to livestock.

Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

28 days

Rates

[field_rates 0](#)

[field_rates 1](#)

-

Restricted Entry Interval

12 hours

Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4-days.

Timings

[when mites first appear](#)