

## **PEARS (INCLUDING ORIENTAL PEAR TREES)**

### General Information

Resistance Management: Because of the inherent risks of resistance development to any product, it is strongly advised that Solera Abamectin Agricultural Miticide/Insecticide be used in a sound resistance management program that includes rotation with other products with different modes of action.

### Limitations, Restrictions, and Exceptions

#### PEARS (INCLUDING ORIENTAL PEAR TREES)

##### Ground Application Only

Horticultural Spray Oil (For best results, use oil with Solera Abamectin Agricultural Miticide/Insecticide).

For Dilute Sprays: 0.25% or 1 gal./A

For Concentrate Sprays: Minimum of 1 gal./A

#### PEARS (INCLUDING ORIENTAL PEAR TREES)

1 The rate of Solera Abamectin Agricultural Miticide/Insecticide per 100 gals. is based on a volume of 400 gals./A dilute spray.

2 To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.

### Application Instructions

- Solera Abamectin Agricultural Miticide/Insecticide may be applied using conventional dilute or concentrate ground sprayers calibrated to deliver sufficient water for thorough coverage. Gallonage of spray will vary with size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for

good spider mite and insect control.

- Always apply in combination with horticultural spray oil (not a dormant oil) as directed. Do not graze treated orchards.

Use Solera Abamectin Agricultural Miticide/Insecticide with a minimum of 0.25% horticultural spray oil (not a dormant oil) in the dilute spray mixture and not less than 1.0 gal. of horticultural spray oil (not a dormant oil) per acre in the final finished spray.

Use Restrictions – Pear

- Do not apply less than 40 gals. of water per acre.
- If a second application is necessary, do not retreat within 21 days of the first application.
- Do not apply by aircraft.
- Do not graze treated orchards.

Method

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

28 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4-days.

Timings

[N.A.](#)