

STRAWBERRIES

General Information

Resistance Management: Because of the inherent risks of resistance development to any product, it is strongly advised that Solera Abamectin Agricultural Miticide/Insecticide be used in a sound resistance management program that includes rotation with other products with different modes of action.

Limitations, Restrictions, and Exceptions

STRAWBERRIES

Application Instructions

- Adjust spray volume and nozzle placement to assure maximum coverage of tops and undersides of leaves.
- Use 16 fl. oz. of Solera Abamectin Agricultural Miticide/Insecticide per acre. Make 2 applications 7-10 days apart when mites first appear.
- Repeat this sequence of application if necessary to maintain control.

Use Restrictions - Strawberries

- For resistance management purposes, do not use in strawberry nurseries.
- Do not apply in less than 100 gals. of water per acre.
- Do not repeat treatment within 21 days of second application.
- Do not apply by aircraft.

Note: Solera Abamectin Agricultural Miticide/Insecticide may be used without any wetting agent. When necessary to improve the wetting of foliage and to smooth out spray deposits, a nonionic surfactant is recommended.

Method

[Spray](#)

Pre-Harvest Interval

3 days

Rates

[field_rates 0](#)

•

Restricted Entry Interval

12 hours

Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4-days.

Timings

[N.A.](#)