

TREE NUTS CROP GROUP (CROP GROUP 14) AND PISTACHIO

General Information

Resistance Management: Because of the inherent risks of resistance development to any product, it is strongly advised that Solera Abamectin Agricultural Miticide/Insecticide be used in a sound resistance management program that includes rotation with other products with different modes of action.

Limitations, Restrictions, and Exceptions

TREE NUTS CROP GROUP (CROP GROUP 14) AND PISTACHIO

Solera Abamectin Agricultural Miticide/Insecticide may be applied using conventional dilute or concentrate sprayers calibrated to deliver sufficient water for thorough coverage. Gallons of spray will vary with size and number of trees per acre and density of foliage. In any case, thorough coverage is essential for good spider mite and insect control.

Instructions

- Always use Solera Abamectin Agricultural Miticide/Insecticide in combination with a horticultural spray oil (not a dormant oil) approved for use on tree nuts.
- Apply Solera Abamectin Agricultural Miticide/Insecticide when spider mites first appear. Residual spider mite control is greater from spray deposits on newer leaves compared to older leaves.
- A second application may be made, if needed, to maintain control. (See Use Restrictions)

1 The rate of Solera Abamectin Agricultural Miticide/Insecticide per 100 gals. is based on a volume of 400 gals./A dilute spray.

2 To determine the amount of product per acre for concentrate sprays, first determine the amount that would be required in a full cover dilute spray. Use the same amount of product per acre in concentrate sprays as would be required for the dilute sprays to the same orchard/grove. This may result in use of less than 10 fl. oz./A on small trees.

Use Restrictions – Tree Nuts Crop Group and Pistachio

- Application Method: Ground application only. Do not apply with aircraft.
- Amount of Water: Do not apply in less than 40 gals. of water per acre.
- Application Interval: If a second application is necessary, do not retreat within 21 days.
- Application per Season: Do not make more than 2 applications of Solera Abamectin Agricultural Miticide/Insecticide or any other foliarly applied abamectin containing product in a growing season.
- Grazing: Do not allow livestock to graze in treated groves/orchards.

Method

[Broadcast/Foliar Ground](#)

Pre-Harvest Interval

21 days

Rates

[field rates 0](#)

[field rates 1](#)

•

Restricted Entry Interval

12 hours

Exception: For grape girdling, cane turning, and tying in grapes, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4-days.

Timings

When spider mites first appear.