

## **APPLICATION TO TURFGRASS (SUPPRESSION)**

### General Information

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### Limitations, Restrictions, and Exceptions

#### APPLICATION TO TURFGRASS:

The active ingredient in GRUBS AWAY has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when application are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation o rainfall to move the active ingredient through the thatch. Do not make applications when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these condition exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

## Application Instructions:

Apply as a broadcast application.

GRUBS AWAY is not effective against mole crickets if adult or large nymphs are present and actively tunneling. A remedial insecticide should be used. Consult your local State Agricultural Experiment station, or State Extension turf specialist for more specific information regarding timing of application.

Note: For optimum control, follow with irrigation or rainfall within 24 hours after application to move the active ingredient through the thatch. On golf courses, irrigate treated areas following application. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

- Refer to label for Spreader Settings.

## RESTRICTIONS:

Do not use GRUBS AWAY on commercial sod farms.

## APPLICATION EQUIPMENT:

Apply GRUBS AWAY uniform over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply GRUBS AWAY. Do not use of spreaders which would apply the material in concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

## Method

[Broadcast Application](#)

## Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

- 

## Timings

[Mole crickets: Prior to or during the peak egg hatch period.](#)

[Chinch bugs: Prior to the hatching of the first instar nymphs.](#)