

# **ASPARAGUS**

## General Information

### PRODUCT INFORMATION

The following directions apply to all uses of DICAMBA 4 DMA. Additional precautions and restrictions will be found in each specific use section.

Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

Do not apply this product through any type of irrigation system.

Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) DICAMBA 4 DMA per application with no more than 2 applications per year.

### MIXING AND APPLICATION

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF DICAMBA 4 DMA. REFER TO INDIVIDUAL AGRICULTURAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

DICAMBA 4 DMA is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (See COMPATIBILITY TEST) should be made prior to tank mixing.

Ground or aerial application equipment, which will give good spray coverage of weed foliage, should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment or 1 to 10 gallons of diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, DICAMBA 4 DMA should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

## BEST STEWARDSHIP PRACTICES

DICAMBA 4 DMA provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading, and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through the soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## SENSITIVE CROP PRECAUTIONS

DICAMBA 4 DMA may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to DICAMBA 4 DMA during their development or growing stage. FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING DICAMBA 4 DMA.

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of DICAMBA 4 DMA with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when air currents may carry spray particles to areas where sensitive crops and plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to

drift out of the target area than fine sprays.

- Use coarse sprays to avoid potential herbicide drift. Select nozzles, which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground application are Delavan Rain-drops, Spraying Systems XR flat fans, or large capacity flood nozzles such as D10, TK10, or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 GPA, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- Do not apply DICAMBA 4 DMA adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.
- To avoid injury to desirable plants, equipment used to apply DICAMBA 4 DMA should be thoroughly cleaned (See PROCEDURE FOR CLEANING SPRAY EQUIPMENT) before reusing to apply any other chemicals.

All crop uses of DICAMBA 4 DMA are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. Tank mix recommendations are for use only in states where the tank mix product and application site are registered.

## BAND TREATMENTS

DICAMBA 4 DMA may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

Refer in the label for tank mix information.

## Limitations, Restrictions, and Exceptions

### ASPARAGUS

FOR USE ONLY IN THE STATES OF CALIFORNIA, OREGON, AND WASHINGTON

Observe all precautions. Read and follow mixing and application instructions.

#### NOTE:

- If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.
- Do not harvest prior to 24 hours after treatment.
- Do not use in the Coachella Valley of California.
- Multiple applications may be made per growing season; however, DO NOT EXCEED a total of 1 pint (1/2 lb. a.i.) of DICAMBA 4 DMA per treated acre per crop year.

#### RATES AND TIMINGS

Apply DICAMBA 4 DMA to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

DICAMBA 4 DMA may be applied in a tank mixture with either 2,4-D or glyphosate herbicide for improved control of noted (\*) weeds. READ AND FOLLOW 2,4-D OR GLYPHOSATE PRODUCT LABELING FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

#### Method

[Broadcast/Foliar Air](#)

[Broadcast/Foliar Ground](#)

#### Rates

[field\\_rates 0](#)

[field\\_rates 1](#)

•

#### Restricted Entry Interval

24 hours

Timings

[Postemergence \(Weed\)](#)